


**Vegetables under Solar PV
2016-17**

Stephen Herbert
herbert@umass.edu

Rationale

- There is a need for sustainable renewable energy sources.
- We suggest solar power as an area of greatest promise for Massachusetts.
- However, traditional ground mounted solar installations on farmland remove arable land from potential agricultural use.

Agricultural Test Site



**UMass Farm South Deerfield
106 Panels ~25 kW**

Solar Array Layout

- Panels were installed (2010) about 7.5ft (2.3m) off the ground.
- There were spaces between panel clusters varying from 2 to 5ft.
- Non-invasive design was used.
- Initial crop evaluation was with pasture and grazing with cattle (2011-2014).











Broccoli, Swiss Chard, Kale, Pepper under shade and unshaded plots transplanted June 7, 2016.







Future crops: Common Bean and Cabbage were planted as second summer crops. Cabbage failed because of the heat and drought in 2016.















Morning Shade Pattern on Test Area prior to Planting
















Initial Conclusions -2016

- All crops, Broccoli, Swiss Chard, Kale, Pepper and Common Bean grew successfully during the drought and heat of the 2016 summer in Massachusetts, however, cabbage failed due to heat stress.
- Leaf temperature was 15°F cooler under the shade of PV panels on clear days contributing to higher yields of shade plots vs. unshaded.
- More shade decrease yield in some crops but not all, and bean had higher yield with more shade possibly related to lower heat stress in mid to late summer.
- Crops were watered daily due to the lack of summer rainfall so moisture availability was not an issue.

Initial Conclusions -2017

- All crops, Broccoli, Swiss Chard, Kale, Pepper (first crops), and Common Bean and Cabbage (second crops) grew successfully during the cooler summer in 2017 in Massachusetts.
- The lack of heat stress on most days in 2017 resulted in higher yields of unshaded plots vs. shaded plots.
- More shade decrease yield in some crops but not all, and the higher yield bean with shade was reversed for unshaded related to lower heat stress in 2017.
- Crops were watered as needed in 2017 which had good summer rainfall.