



A Pictorial Guide to Hedgerow Plants for Beneficial Insects

By Omar Rodriguez and Rex Dufour, NCAT Agriculture Specialists
Published Oct. 2017
©NCAT
IP547

This publication provides a straightforward pictorial guide to several of the most beneficial hedgerow plant species used in farmscaping for native pollinators and insect predators and parasites in California. It provides plant names, bloom times, heights, and descriptions that note considerations for selection and establishment.

Contents

- Introduction.....1
- Hedgerow Plants.....2
- Additional Considerations 10
- Further Resources 10



Introduction

When selecting the appropriate perennial plants to use in your hedgerow, it is important to define the benefits you would like to see as a result of adding hedgerows to your farm. This informational piece highlights a collection of native California species (with the exception of lavender) that are both drought tolerant and highly attractive to beneficial insects. In addition to attracting beneficial insects (predators, parasites, and pollinators), hedgerows can serve a number of other functions that include providing overwintering sites for desirable insect species, wind breaks, dust barriers, pesticide barriers, bird nesting and perches, carbon sequestration, stream-bank stabilization or revegetation, and shade for stream cooling and fish habitat, not to mention simply adding some practical beauty to an agricultural landscape.

Consider the effects that each hedgerow species' size and bloom time will have on your farm. The impacts of planting hedgerows are heavily on the positive side, but it's important to understand the exceptions. For example, toyon, which is explored in this publication, is a useful species that attracts many beneficial insects and birds. However, it is also susceptible to fire blight, so it would be inappropriate for a border planting for an apple or pear orchard.

A biodiverse hedgerow provides farm resiliency to the pressures that pests impose, which include not only destructive habits such as eating or boring into plants. Pests are also carriers and transmitters of bacteria, viruses, and fungi that cause disease in the plants we grow. Creating habitat for beneficial insects is one step in better managing and supporting these "mini-livestock," which can provide many benefits to your farm.



Funding for this publication was provided in part by the National Institute of Food and Agriculture, Award 2015-70017-22868 of the Beginning Farmer and Rancher Development Program (BFRDP), part of a subcontract administered by UC Berkeley, PI's Jennifer Sowerwine and Christy Getz.

ATTRA (www.attra.ncat.org) is a program of the National Center for Appropriate Technology (NCAT). The program is funded through a cooperative agreement with the United States Department of Agriculture's Rural Business-Cooperative Service. Visit the NCAT website (www.ncat.org) for more information on our other sustainable agriculture and energy projects.



Hedgerow Plants

California Buckwheat

Scientific name: *Eriogonum fasciculatum*

Mature height: 2 to 4 feet

Bloom time: April to September

Notes: Very drought tolerant, an important food source for many native bees and beneficial insects, including syrphid flies (also known as flower flies or hover flies), predatory wasps, pirate bugs, tachinid flies, and lady beetles. Full sun to partial shade. Blooms are creamy white, last through the summer, and turn an attractive rust color when they dry down.



Assassin bugs are generalist predators with a wide host range. Photo: Rex Dufour, NCAT



Tachinid fly on California buckwheat. Tachinid larvae are parasites of stinkbugs and other insects. Photo: Rex Dufour, NCAT



California buckwheat in bloom. Photo: Rex Dufour, NCAT

California Fuchsias

Scientific name: *Epilobium canum*

Mature height: 1 to 4 feet

Bloom time: July to October

Notes: May need help establishing through its first summer. Highly tolerant in a variety of substrates and one of the most attractive flowers to hummingbirds, which consume a variety of insects, including aphids. The bright-red, tubular flowers provide a welcome visual relief during the hottest, driest, and dustiest part of the season, as well as a nectar and pollen resource for beneficials.



The bright red-orange flowers can offer beneficials a pollen and nectar source during the drier, dustier parts of the season, August and September. Photo: Rex Dufour, NCAT



California fuchsia. Photo: Rex Dufour, NCAT



Close up of California fuchsia. Photo: Rex Dufour, NCAT



Honey bee gathering nectar and pollen from early spring ceanothus bloom. Photo: Rex Dufour, NCAT

Ceanothus/California Lilac

Scientific name: *Ceanothus* sp. (many varieties exist)

Mature height: 2 to 20 feet (depends on variety)

Bloom time: March to May

Notes: These plants are nitrogen fixers and occur naturally on dry slopes across North America and as far south as Guatemala. Ceanothus does not require summer irrigation after it has been established. Its blue flowers attract many pollinators early in the season, and its foliage maintains a dense green color all year long. Comes in tall and prostrate cultivars. Deer like to browse on this plant.



Ceanothus in full bloom. Photo: Rex Dufour, NCAT



Syrphid fly on ceanothus flower. Syrphid larvae are aphid predators. Photo: Rex Dufour, NCAT



Cleveland sage in bloom. It will spread nicely to cover ground and smother weeds. Photo: Rex Dufour, NCAT

Cleveland Sage

Scientific name: *Salvia clevelandii*

Mature height: 3 feet

Bloom time: April to August

Notes: Has attractive flowers, inviting to several bee species, butterflies, and hummingbirds. Leaves are very aromatic. Prefers good drainage; can tolerate full sun. Watering in early stages will help it establish, but it does well in drought conditions.



Carpenter bee getting nectar from Cleveland sage. Photo: Rex Dufour, NCAT



Lady bird beetles resting on Cleveland sage leaves in early spring. Photo: Rex Dufour, NCAT

Coffee Berry

Scientific name: *Frangula californica*

Mature height: 6 to 15 feet

Bloom time: April to June

Notes: An insect-pollinated plant that attracts a large range of beneficial insects: honey bee, native bees, and flies. It has a relatively short bloom period.



Coffee berries ripening. Photo: Rex Dufour, NCAT



Coffee berry flowers and berries with lady bird beetle. Photo: Rex Dufour, NCAT



Green lacewing adult feeding on coffee berry nectar and pollen. Lacewing larvae are voracious predators of aphids and other soft-bodied insects (mealybugs, small caterpillars, psyllids). Photo: Rex Dufour, NCAT

Coyote Bush

Scientific name: *Baccharis pilularis*

Mature height: 6 to 12 feet

Bloom time: November to February

Notes: Beneficial insectary and wildlife habitat, which hosts many insects even when not in bloom. Very good winter pollen/nectar source. Has male and female plants, and female flowers develop “fluffy” blooms once they develop seeds. Choose between prostrate (up to 3.5 feet high) and non-prostrate (12 feet high) varieties.



Lady bird beetle on female coyote bush flower. Photo: Rex Dufour, NCAT



Coyote bush in bloom. Photo: Rex Dufour, NCAT



Tachinid fly feeding at male coyote bush flower. Photo: Rex Dufour, NCAT



Praying mantis egg mass in the interior of deer grass bunch. Several dozen small praying mantids will emerge from this egg mass. Photo: Rex Dufour, NCAT



Lady bird beetles overwintering in deer grass. Groups of beetles emit an aggregation pheromone that attracts additional lady bird beetles. Photo: Rex Dufour, NCAT



Elderberry buds and blooms. Photo: Rex Dufour, NCAT



Elderberry in a hedgerow. Ideal as windbreak, dust barrier, and bird habitat, but may need some pruning of multiple stem base. Photo: Rex Dufour, NCAT

Deergrass

Scientific name: *Muhlenbergia rigens*

Mature height: 4 to 5 feet and 4 to 6 feet wide

Bloom time: May to September

Notes: Clumping grass; interior is good overwintering habitat for ladybird beetles, and seed spikes are good resting places for damsel and dragon flies. Native to most of California, Texas, and Mexico. Resilient to different soil types. It does best with full sun exposure but will tolerate some shade and is drought tolerant. Due to its abundant yield of seed, it is a great host plant for birds, as well as beneficial insects.



Deergrass in a farm hedgerow. Photo: Rex Dufour, NCAT

Elderberry

Scientific name: *Sambucus nigra*

Mature height: 6 to 14 feet

Bloom time: May to August

Notes: Can be found in many riparian habitats throughout California and New Mexico. It grows quickly and will need regular watering in drier regions, about once every two weeks during the summer. Its flowers attract many beneficials. Flowers give way to fall berries that bring in many species of songbirds. Prefers full sun.



Elderberry fruit, attractive to birds, and some jam makers. Photo: Rex Dufour, NCAT

Flannel Bush

Scientific name: *Fremontodendron californicum*

Mature height: 8 to 25 feet

Bloom time: April to June

Notes: Has many large, bright-yellow blooms that attract bees, wasps, and flies of many species. Grows rapidly and is drought tolerant.



Flannel bush in full bloom. Photo: Rex Dufour, NCAT



The large flowers with easily accessed pollen are very attractive to honey bees. Photo: Rex Dufour, NCAT



Flannel bush flowers are large (3-4" across) and eye catching. Photo: Rex Dufour, NCAT

Hollyleaf Cherry

Scientific name: *Prunus ilicifolia*

Mature Height: 8 to 20 feet

Bloom Time: April to May

Notes: Very attractive to native bees and parasitic flies. Can grow quite large under favorable growth conditions; prune to shape.



A holly leaf cherry tree in full bloom represents a gold mine of pollen and nectar for beneficial insects. Photo: Rex Dufour, NCAT



Syrphid fly on hollyleaf cherry blossom. Although the bloom time is a relatively short two weeks for an individual plant, when these are in bloom, the buzz of thousands of insects can be heard. Photo: Rex Dufour, NCAT



Holly leaf cherry fruit. Photo: Rex Dufour, NCAT



Lavender in bloom. Although not native to California, it does very well in this Mediterranean climate. Photo: Rex Dufour, NCAT



Lavender in bloom. Many types of bees are attracted to these flowers. Photo: Rex Dufour, NCAT

Lavender

Scientific name: *Lavandula sp.*

Mature height: 1 to 4 feet

Bloom time: June to August

Notes: Lavender is not native to California, but it is a favorite of both native bees and honeybees. This woody shrub is drought tolerant and flourishes in dry, well-drained soils.



Lavender is popular with bees and some beekeepers believe lavender nectar provides medicinal benefits to their bees. Photo: Rex Dufour, NCAT



Narrowleaf milkweed in bloom. Planting these will provide habitat for the monarch butterfly. Photo: Rex Dufour, NCAT

Narrowleaf Milkweed

Scientific name: *Asclepias fascicularis*

Mature height: 2 to 4 feet

Bloom time: June to September

Notes: This plant is host for the Monarch butterfly larvae and the bright-yellow oleander aphid, which does not attack commercial food crops in California. The oleander aphids provide alternate prey for syrphid fly larvae, green lacewing larvae, ladybird beetles, and the aphid is an alternate host for several parasitic wasps.



Close up of narrowleaf milkweed flowers in bloom. The yellow dots on the stems are oleander aphids, alternate prey for aphid parasites and predators. Photo: Rex Dufour, NCAT



Monarch butterfly larva, chrysalis and oleander aphids on narrow leaf milkweed. Monarch butterfly populations have recently decreased dramatically and farmers can help provide habitat for these creatures. Photo: Rex Dufour, NCAT

Quailbush/Saltbush

Scientific name: *Atriplex nummularia*

Mature height: Up to 10 feet

Bloom Time: August to October

Notes: Saltbush is a drought-tolerant, native shrub in the daisy family that can be found throughout the western United States. Its services as a native pollinator attractant are valuable in that it flowers at a time of year when most other plants are dormant or have finished blooming. Additionally, it is an important nectar source for migrating monarch butterflies. Saltbush is tolerant of saline soils and competes well against weeds but can be somewhat aggressive in its growth.



Syrphid fly on Saltbush flower. Syrphid larvae are aphid predators. Photo: Rex Dufour, NCAT



Honey bee gathering nectar and pollen from early fall Saltbush bloom. Photo: Rex Dufour, NCAT



Saltbush in full bloom. Photo: Rex Dufour, NCAT

Toyon

Scientific name: *Heteromeles arbutifolia*

Mature height: 8 to 20 feet and equally as wide

Bloom time: June to August

Notes: An evergreen shrub, also known as Christmas berry. White flower bunches attract bees; bright-red berries attract birds. It has a profuse root system that can help with erosion control and stabilizing hillsides. Susceptible to fireblight, so don't plant with apples or pears.



Toyon in bloom. Photo: Rex Dufour, NCAT



Toyon flowers and buds. Honey bees and many beneficials are attracted to toyon. Photo: Rex Dufour, NCAT



December toyon with red berries next to deer grass. It's clear why toyon is also called "Christmas berry." Photo: Rex Dufour, NCAT



Redbud in bloom in early spring. Note the multiple stems. Photo: Rex Dufour, NCAT

Western Redbud

Scientific name: *Cercis orbiculata*

Mature height: 7 to 20 feet

Bloom time: March to May

Notes: Native to parts of California, Arizona, and Utah. It is drought tolerant once established but may require regular watering in desert environments. Full sun to partial shade, and it does best in well-drained soil. Attractive to many pollinators and beneficial insects and generally pollinated by bumble bees and mason bees. Leafcutter bees also like to use its soft new leaves to line their nests. Redbud also re-sprouts vigorously after fires.



Redbud leaves. Photo: Rex Dufour, NCAT



Red bud is in the legume family, and seeds can be collected for planting from the pods, shown here. Photo: Rex Dufour, NCAT



Yarrow plant in bloom. Photo: Rex Dufour, NCAT

Yarrow

Scientific name: *Achillea millefolium*

Mature height: 1.5 to 3 feet

Bloom time: April to September

Notes: Yarrow, a hardy perennial, is attractive to a number of pollinators and predatory species like lady beetles and syrphid flies. There are pink (salmon-colored), yellow, and white varieties. It is very resilient, water needs are minimal (it doesn't like wet feet), and it will establish itself in a variety of soil types.



Native bee on yarrow. Photo: Rex Dufour, NCAT



Native bees, which are often more effective pollinators than honey bees, are attracted to yarrow. Photo: Rex Dufour, NCAT

Additional Considerations

Potential hedgerow plants are available in a wide range of sizes and types from local nurseries specializing in native plants (see Further Resources, below). Ideally, hedgerows should be designed to include plants that bloom in different seasons to provide pollinators and other beneficial insects with food sources throughout the growing season. Structural diversity

is also important in order to support a diverse ecology of beneficials. Finally, remember that hedgerow site preparation is important to successfully establishing a hedgerow. Weed competition needs to be minimized, either through solarizing, use of weed barriers, regular (and heavy) mulching, hand weeding, or use of herbicides.

Further Resources

Suppliers

For a listing of native plant nurseries in California, go to: www.plantnative.org/nd_ca.htm

Publications

Beneficial Insect Habitat Assessment Form and Guide. 2015. By The Xerces Society for Invertebrate Conservation. www.xerces.org/wp-content/uploads/2015/07/HAG_BeneficialInsects_June2015_web.pdf

Companion Planting & Botanical Pesticides: Concepts and Resources. 2016. By George Kuepper, Mardi Dodson, and Justin Duncan. National Center for Appropriate Technology. Publication IP125. www.attra.ncat.org/attra-pub/summaries/summary.php?pub=72

Conservation Buffers in Organic Systems. 2014. By National Center for Appropriate Technology, Oregon Tilth, and The Xerces Society for Invertebrate Conservation. National Center for Appropriate Technology. Publication IP470. www.attra.ncat.org/attra-pub/summaries/summary.php?pub=464

Establishing Hedgerows on Farms in California. 2010. By University of California Agriculture and Natural Resources. Publication 8390. <http://ucfoodsafety.ucdavis.edu/files/26499.pdf>

Farming With Native Beneficial Insects. 2014. By Eric Lee-Mäder, Jennifer Hopwood, Mace Vaughan, Scott Hoffman Black, and Lora Morandin. The Xerces Society for Invertebrate Conservation. <https://xerces.org/farming-with-native-beneficial-insects>

Farmscaping to Enhance Biological Control. 2000. By Rex Dufour. National Center for Appropriate Technology. Publication CT065. <http://extension.oregonstate.edu/sorec/sites/default/files/farmscaping.pdf>

Hedgerows for California Agriculture: A Resource Guide. 2004. By Sam Earnshaw. Community Alliance with Family Farmers (CAFF). www.caff.org/wp-content/uploads/2010/07/Hedgerow_manual.pdf

Pollinator Plants, California. 2014. By The Xerces Society for Invertebrate Conservation. www.xerces.org/wp-content/uploads/2014/09/CaliforniaPlantList_web.pdf

Online Resources

Native Plant Lists by Region. California Native Plant Society. www.cnps.org/cnps/grownative/lists.php

This publication was developed with support from NIFA/BFRDP grant number 201570001722868.

**A Pictorial Guide to Hedgerow Plants for
Beneficial Insects**

By Omar Rodriguez and Rex Dufour,

NCAT Agriculture Specialists

Published October 2017 © NCAT

Tracy Mumma, Editor • Amy Smith, Production

This publication is available on the Web at:

www.attra.ncat.org

IP547

Slot 570

Version 101117